



Aceite de la palma de seje *Oenocarpus bataua* Mart. por su calidad nutricional puede contribuir a la conservación y uso sostenible de los bosques de galería en la Orinoquia Colombiana [

2013

text (article)

Analítica

The riparian forest on eastern plains of Colombia, many of them rich on seje palm (*Oenocarpus bataua* Mart.), have been meaning as nutritional support for ancestral communities of the region. The extraction of seje Oil was done in the Natural Reserve the Unamas, San Martin county, 300 a.s.l., Meta department, Colombia. A three riparian forest of 900 hectares was used for the study. The seje palm and its oil extraction illustrate the potential forest sustainable use, with seje oil, which is similar or superior to olive oil, edible for human consumption and strategic potential use for animal nutrition. A density of palms in the riparian forest was established; soil quality and protocols for the seje oil extraction were evaluated, trying to improve the traditional oil extraction protocol, management the temperature during the process. For the traditional protocol 1 the maximum oil extraction was 2,5% while with protocol 2 it was of 4,26% improving the oil extraction on 70%. The average palm density was 187,7 ha-1 where 67% of the palms were on production (125,7 palms ha-1) and 33% were unproductive on development (62 palms ha-1). The soil were the seje palm is produced is acid, low in phosphorus and nitrogen content, and high aluminium and iron concentration, but the forest productivity seems to be linked to nutrient cycling by trash and microorganism on the superficial soil layer. The seje oil composition was not affected by fruit ripeness but the fruit ripe produced more oil. The seje oil is rich on oleic acid similar to olive oil, with lower linoleic acid but higher linolenic acid content. The seje oil product of high nutritional value, a product of riparian forest, should contribute to forest conservation and its sustainable use

The riparian forest on eastern plains of Colombia, many of them rich on seje palm (*Oenocarpus bataua* Mart.), have been meaning as nutritional support for ancestral communities of the region. The extraction of seje Oil was done in the Natural Reserve the Unamas, San Martin county, 300 a.s.l., Meta department, Colombia. A three riparian forest of 900 hectares was used for the study. The seje palm and its oil extraction illustrate the potential forest sustainable use, with seje oil, which is similar or superior to olive oil, edible for human consumption and strategic potential use for animal nutrition. A density of palms in the riparian forest was established; soil quality and protocols for the seje oil extraction were evaluated, trying to improve the

traditional oil extraction protocol, management the temperature during the process. For the traditional protocol 1 the maximum oil extraction was 2,5% while with protocol 2 it was of 4,26% improving the oil extraction on 70%. The average palm density was 187,7 ha-1 where 67% of the palms were on production (125,7 palms ha-1) and 33% were unproductive on development (62 palms ha-1). The soil where the seje palm is produced is acid, low in phosphorus and nitrogen content, and high aluminium and iron concentration, but the forest productivity seems to be linked to nutrient cycling by trash and microorganism on the superficial soil layer. The seje oil composition was not affected by fruit ripeness but the fruit ripe produced more oil. The seje oil is rich on oleic acid similar to olive oil, with lower linoleic acid but higher linolenic acid content. The seje oil product of high nutritional value, a product of riparian forest, should contribute to forest conservation and its sustainable use.

The riparian forest on eastern plains of Colombia, many of them rich on seje palm (*Oenocarpus bataua* Mart.), have been meaning as nutritional support for ancestral communities of the region. The extraction of seje Oil was done in the Natural Reserve the Unamas, San Martin county, 300 a.s.l., Meta department, Colombia. A three riparian forest of 900 hectares was used for the study. The seje palm and its oil extraction illustrate the potential forest sustainable use, with seje oil, which is similar or superior to olive oil, edible for human consumption and strategic potential use for animal nutrition. A density of palms in the riparian forest was established; soil quality and protocols for the seje oil extraction were evaluated, trying to improve the traditional oil extraction protocol, management the temperature during the process. For the traditional protocol 1 the maximum oil extraction was 2,5% while with protocol 2 it was of 4,26% improving the oil extraction on 70%. The average palm density was 187,7 ha-1 where 67% of the palms were on production (125,7 palms ha-1) and 33% were unproductive on development (62 palms ha-1). The soil where the seje palm is produced is acid, low in phosphorus and nitrogen content, and high aluminium and iron concentration, but the forest productivity seems to be linked to nutrient cycling by trash and microorganism on the superficial soil layer. The seje oil composition was not affected by fruit ripeness but the fruit ripe produced more oil. The seje oil is rich on oleic acid similar to olive oil, with lower linoleic acid but higher linolenic acid content. The seje oil product of high nutritional value, a product of riparian forest, should contribute to forest conservation and its sustainable use.

<https://rebiunoda.pro.baratznet.cloud:28443/OpacDiscovery/public/catalog/detail/b2FpOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzI5Mjk3ODg>

Título: Aceite de la palma de seje *Oenocarpus bataua* Mart. por su calidad nutricional puede contribuir a la conservación y uso sostenible de los bosques de galería en la Orinoquia Colombiana [electronic resource]

Editorial: 2013

Tipo Audiovisual: Aceite de seje bosque de galería orinocense ácido linoléico ácido linolénico ácido oleico *Oenocarpus bataua* Mart palma de seje Seje oil riparian orinoquia forest oleic acid linoleic acid linolenic acid *Oenocarpus bataua* Mart seje palm Óleo Seje Floresta de galería linoléico linolénico *Oenocarpus bataua* Mart oléico e Palma Seje

Documento fuente: Orinoquía, ISSN 0121-3709, Vol. 17, N°. 2, 2013, pags. 215-229

Nota general: application/pdf

Restricciones de acceso: Open access content. Open access content star

Condiciones de uso y reproducción: LICENCIA DE USO: Los documentos a texto completo incluidos en Dialnet son de acceso libre y propiedad de sus autores y/o editores. Por tanto, cualquier acto de reproducción, distribución, comunicación pública y/o transformación total o parcial requiere el consentimiento expreso y escrito de aquéllos. Cualquier enlace al texto completo de estos documentos deberá hacerse a través de la URL oficial de éstos en Dialnet. Más información: <https://dialnet.unirioja.es/info/derechosOAI> | INTELLECTUAL PROPERTY RIGHTS STATEMENT: Full text documents hosted by Dialnet are protected by copyright and/or related rights. This digital object is accessible without charge, but its use is subject to the licensing conditions set by its authors or editors. Unless expressly stated otherwise in the licensing conditions, you are free to linking, browsing, printing and making a copy for your own personal purposes. All other acts of reproduction and communication to the public are subject to the licensing conditions expressed by editors and authors and require consent from them. Any link to this document should be made using its official URL in Dialnet. More info: <https://dialnet.unirioja.es/info/derechosOAI>

Lengua: Spanish

Enlace a fuente de información: Orinoquía, ISSN 0121-3709, Vol. 17, Nº. 2, 2013, pags. 215-229

Baratz Innovación Documental

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- informa@baratz.es